




## SCIENTIFIC PAPER

### A FAUNISTIC STUDY ON SOME FAMILIES OF CHALCIDOIDEA (HYMENOPTERA) FROM GOLESTAN PROVINCE AND VICINITY, NORTHERN IRAN

Hassan Ghahari<sup>1</sup>

Svetlana N. Myartseva<sup>2</sup>

Enrique Ruíz-Cancino<sup>2</sup> 

<sup>1</sup> Department of Plant Protection, Yadegar – e-Imam Khomeini (RAH) Branch Islamic Azad University.  
hghahari@yahoo.com

<sup>2</sup> Facultad de Ingeniería y Ciencias Universidad Autónoma de Tamaulipas  
smyartse@uat.edu.mx

 [eruz@uat.edu.mx](mailto:eruz@uat.edu.mx)

<sup>1</sup> Sciences & Research Branch, Hesarak, Tehran, I. R. Iran.

<sup>2</sup> Universidad Autónoma de Tamaulipas, C. P. 87000 Ciudad Victoria, Tamaulipas, México.

*Folia Entomológica Mexicana (nueva serie)*, 2(1): 10–15, 2016.

Recibido: 27 de octubre 2015

Aceptado: 13 de marzo 2016

Publicado en línea: 30 de abril 2016

**A FAUNISTIC STUDY ON SOME FAMILIES OF CHALCIDOIDEA  
(HYMENOPTERA) FROM GOLESTAN PROVINCE AND VICINITY, NORTHERN  
IRAN**

**Estudio Faunístico de algunas familias de CHALCIDOIDEA (HYMENOPTERA) de la  
provincia de Golestán y sus alrededores, al norte de Irán**

Hassan Ghahari<sup>1</sup>, Svetlana N. Myartseva<sup>2</sup> and Enrique Ruíz-Cancino<sup>2\*</sup>

<sup>1</sup>Islamic Azad University, Tehran, Iran.

<sup>2</sup>Universidad Autónoma de Tamaulipas.

\*Corresponding author [eruiz@uat.edu.mx](mailto:eruiz@uat.edu.mx)

---

**ABSTRACT.** The fauna of seven families of Chalcidoidea from Golestan Province and vicinity in northern Iran was studied. The families determined were: Chalcididae (2 species), Eucharitidae (1), Eulophidae (2), Eurytomidae (8), Ormyridae (1), Perilampidae (2), and Torymidae (3). Twelve species are new records for the fauna of Iran.

**Key words:** Fauna, new records, parasitoids.

**RESUMEN.** Se estudió la fauna de siete familias de Chalcidoidea de la Provincia de Golestán y sus alrededores, en el norte de Irán. Las familias determinadas fueron: Chalcididae (2 especies), Eucharitidae (1), Eulophidae (2), Eurytomidae (8), Ormyridae (1), Perilampidae (2) y Torymidae (3). Doce especies son nuevos registros de la fauna de Irán.

**Palabras clave:** Fauna, nuevos registros, parasitoides.

---

## INTRODUCTION

Chalcidoidea includes 19 families of parasitic hymenopterans that range in size from 0.1 mm to 45 mm. These wasps parasitize a wide variety of immature arthropods: 13 orders of Insecta, two orders of Arachnida, and one family of Nematoda (Grissel and Schauff, 1997). Members of this superfamily are cosmopolitan in distribution and habitat utilization (Noyes and Hayat, 1984; LaSalle y Gauld, 1991). Chalcidoids have a greater range of biological diversity than the species of other parasitoids. Most species are parasitoids, but more than 80 species in some families especially Eurytomidae and Torymidae are phytophagous (Gauld and Bolton, 1988; Noyes y Valentine, 1989). Chalcidoidea is the most important successful group used in applied biological control. They have been used widely in

successful biological control programs of insect pests, four of the six most successful hymenopteran families utilized belong to Chalcidoidea (Noyes and Hayat, 1994; Godfray, 1994).

Golestan province (36.8393° N 54.4444° W) is located in the north of Iran and south of the Caspian Sea, geographically it is divided into two sections, the plains and the mountains of the Alborz range, in the eastern Alborz section. The mountains have a northeasterly aspect and gradually decrease in height; the highest point of the province is Shavar with 3945 m above sea level; climate of Golestan is temperate most of the year (Hassan *et al.*, 1993). The aim of this research was a faunistic survey on the families Chalcididae, Eucharitidae, Eulophidae, Eurytomidae, Ormyridae, Perilampidae, and Torymidae from Golestan province, Northern Iran.

## MATERIAL AND METHODS

The fauna of some families of Iranian Chalcidoidea was studied in some regions of Golestan province and vicinity (Northern Iran). The specimens were collected by means of sweeping nets and Malaise traps. In addition, some parasitoids were obtained through the rearing of their hosts in optimum condition ( $27 \pm 1^\circ\text{C}$ ,  $70 \pm 5\%$  RH, 16: 8 L: D) in an incubator for emergence of parasitoids. Taxonomical determinations were done using identification keys (e.g. Peck *et al.*, 1964; Boucek, 1988; Sharma, 1988; Grissell and Schauff, 1997). In this paper, classification, nomenclature and distribution data were taken from Noyes (2014).

## RESULTS

In this research, a total of 19 species belong to seven families of Chalcidoidea were collected and identified from Golestan province and vicinity, northern Iran. Twelve species, *Brachymeria moerens* (Ruschka) (Chalcididae), *Eucharis adscendens* (Fabricius) (Eucharitidae), *Elasmus flabellatus* (Fonscolombe) (Eulophidae), *Bruchophagus coluteae* Boucek, *Eurytoma infracta* Mayr, *E. oophaga* Silvestri, *Tetramesa agrostidis* (Howard), *T. fumipennis* Walker (Eurytomidae), *Chrysolampus prominens* (Ruschka), *C. rufitarsis* (Förster) (Perilampidae), *Pseudotorymus sapphyrinus* (Fonscolombe), and *Torymus nitidulus* (Walker) (Torymidae) are new records for the fauna of Iran. The list of species with distributional data is given below alphabetically.

Family Chalcididae Latreille, 1817  
Genus *Brachymeria* Westwood, 1829

*Brachymeria moerens* (Ruschka, 1922)

**Material examined:** Golestan province, Minudasht,  $37^\circ 10' \text{N}$   $55^\circ 30' \text{E}$ , 1, October 2012. **New record for Iran.**

**Distribution outside Iran:** Austria, Bosnia-Herzegovina, Bulgaria, Caucasus, Czech Republic, France, Germany, Hungary, Italy,

Macedonia, Moldova, Slovakia, Transcaucasus, Turkey, Ukraine, and former Yugoslavia.

*Brachymeria obtusata* (Förster, 1859)

**Material examined:** Golestan province, Kordkoy,  $36^\circ 41' \text{N}$   $54^\circ 12' \text{E}$ , 3, 1, August 2009; Golestan National Park, Yakhtikalan (1887 m), 2, June 2012; Mazandaran province, Babol,  $36^\circ 30' \text{N}$   $52^\circ 35' \text{E}$ , 2, June 2014.

**Distribution outside Iran:** Albania, Austria, Croatia, Cyprus, Czech Republic, Egypt, France, Germany, Greece, Hungary, Iraq, Israel, Italy, Japan, Moldova, Netherlands, North Africa, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Transcaucasus, Turkey, Turkmenistan, Ukraine, former USSR, and former Yugoslavia.

Family Eucharitidae Walker, 1846  
Genus *Eucharis* Latreille, 1804

*Eucharis adscendens* (Fabricius, 1787)

**Material examined:** Golestan province, Kordkoy,  $36^\circ 41' \text{N}$   $54^\circ 12' \text{E}$ , 3, 1, August 2009. **New record for Iran.**

**Distribution outside Iran:** Austria, Belarus, Bosnia-Herzegovina, Croatia, Czech Republic, Germany, Hungary, Italy, Kazakhstan, Lithuania, Moldova, Poland, Romania, Russia, Serbia, Siberia, Slovakia, Transcaucasus, Turkey, Ukraine, United Kingdom (UK), and former USSR.

Family Eulophidae Westwood, 1829  
Genus *Elachertus* Spinola, 1811

*Elachertus charondas* (Walker, 1839)

**Material examined:** Golestan province, Gorgan,  $36^\circ 50' \text{N}$   $54^\circ 30' \text{E}$ , 3, July 2011; Mazandaran province, Sari,  $36^\circ 30' \text{N}$   $53^\circ 30' \text{E}$ , 1, April 2013.

**Distribution outside Iran:** Brazil, China, Czech Republic, Germany, Hungary, Italy, Japan, Moldova, Russia, Slovakia, Spain, Sweden, Taiwan, Turkmenistan, UK, and Yemen.

Genus *Elasmus* Westwood, 1833

*Elasmus flabellatus* (Fonscolombe, 1832)

**Material examined:** Golestan province, Gonbad, 37° 30' N 55° 00' E, 2 , September 2013. **New record for Iran.**

**Distribution outside Iran:** Austria, Bosnia-Herzegovina, Bulgaria, Croatia, Czech Republic, France, Germany, Greece, Hungary, Italy, Lebanon, Moldova, Morocco, Netherlands, Portugal, Romania, Russia, Serbia, Slovakia, Spain, Sweden, Turkey, UK, former USSR, and former Yugoslavia.

Family Eurytomidae Walker, 1832

Genus *Bruchophagus* Ashmead, 1888

*Bruchophagus coluteae* Boucek, 1954

**Material examined:** Golestan province, Minudasht, 37° 10' N 55° 30' E, 1 , October 2012. **New record for Iran.**

**Distribution outside Iran:** Bulgaria, Central Asia, Czech Republic, Moldova, Slovakia, Turkey, former USSR, and Uzbekistan.

Genus *Eurytoma* Illiger, 1807

*Eurytoma collaris* Walker, 1832

**Material examined:** Golestan province, Kalaleh, 37° 43' N 55° 49' E, 1 , 1 , October 2012; Northern Khorasan province, Bojnord, 37° 35' N 57° 20' E, 1 , May 2011.

**Distribution outside Iran:** Bulgaria, Czech Republic, France, Germany, Hungary, Kazakhstan, Slovakia, Sweden, Turkey, and UK.

*Eurytoma infracta* Mayr, 1904

**Material examined:** Golestan National Park, Dasht-e-Mirza-Bayloo (1589 m), 2 , 1 , June 2012. **New record for Iran.**

**Distribution outside Iran:** Caucasus, Croatia, France, Jordan, Romania, Spain, Turkey, Ukraine, Central Asia, and former Yugoslavia.

*Eurytoma oophaga* Silvestri, 1920

**Material examined:** Golestan province, Gorgan, 36° 50' N 54° 30' E, 1 , 1 , July 2011. **New record for Iran.**

**Distribution outside Iran:** Caucasus, France, Germany, Italy, Romania, Serbia, Transcaucasus, and Turkey.

Genus *Macrorileyia* Ashmead, 1900

*Macrorileyia inopinata* Silvestri, 1920

**Material examined:** Northern Khorasan province, Bojnord, 37° 35' N 57° 20' E, 1 , May 2011; Golestan province, Gonbad, 37° 30' N 55° 00' E, 1 , September 2013.

**Distribution outside Iran:** Bulgaria, Caucasus, Central Asia, Croatia, Czech Republic, Georgia, Hungary, Italy, Kazakhstan, Mexico, Moldova, North Africa, Romania, Serbia, Slovakia, Spain, Transcaucasus, Tunisia, Turkey, and United States of America (USA).

Genus *Tetramesa* Walker, 1848

*Tetramesa agrostidis* (Howard, 1896)

**Material examined:** Golestan province, Kordkoy, 36° 41' N 54° 12' E, 1 , August 2009; Northern Khorasan province, Bojnord, 37° 35' N 57° 20' E, 1 , May 2011. **New record for Iran.**

**Distribution outside Iran:** Bulgaria, Germany, Kazakhstan, Turkey, USA, and former USSR.

*Tetramesa fulvicollis* Walker, 1832

**Material examined:** Golestan province, Minudasht, 37° 10' N 55° 30' E, 1 , October 2012.

**Distribution outside Iran:** Bulgaria, Croatia, Czech Republic, France, Germany, Hungary, Montenegro, Netherlands, Romania, Slovakia, Sweden, Turkey, UK, and former USSR.

*Tetramesa fumipennis* Walker, 1832

**Material examined:** Golestan National Park, Yakhtikalan (1887 m), 2 , June 2012. **New record for Iran.**

**Distribution outside Iran:** Caucasus, Germany, Hungary, Romania, Sweden, Turkey, UK, and former USSR.

Family Ormyridae Förster, 1856  
Genus *Ormyrus* Westwood, 1832

*Ormyrus diffinis* (Fonscolombe, 1832)

**Material examined:** Northern Khorasan province, Bojnord, 37° 35' N 57° 20' E, 2 , May 2011. Golestan province, Kalaleh, 37° 43' N 55° 49' E, 2 , 1 , October 2012. Mazandaran province, Babol, 36° 30' N 52° 35' E, 1 , June 2014.

**Distribution outside Iran:** Bulgaria, Caucasus, Croatia, Czech Republic, France, Germany, Hungary, Israel, Kazakhstan, Macedonia, Moldova, Montenegro, Russia, Serbia, Slovakia, Spain, Tadjikistan, Transcaucasus, Turkey, Turkmenistan, Ukraine, former USSR, and former Yugoslavia.

Family Perilampidae Förster, 1856  
Genus *Chrysolampus* Spinola, 1811

*Chrysolampus prominens* (Ruschka, 1924)

**Material examined:** Golestan province, Gorgan, 36° 50' N 54° 30' E, 1 , July 2011. **New record for Iran.**

**Distribution outside Iran:** Hungary, Italy, Kazakhstan, Romania, Slovenia, Spain, and Turkmenistan.

*Chrysolampus rufitarsis* (Förster, 1859)

**Material examined:** Golestan National Park, Dasht-e-Mirza-Bayloo (1589 m), 1 , 1 , June 2012. **New record for Iran.**

**Distribution outside Iran:** Austria, Belgium, Canary Islands, Czech Republic, Germany, Hungary, Kazakhstan, Netherlands, Slovakia, Spain, and UK.

Family Torymidae Walker, 1833  
Genus *Pseudotorymus* Masi, 1921

*Pseudotorymus sapphyrinus* (Fonscolombe, 1832)

**Material examined:** Golestan province, Gorgan, 36° 50' N 54° 30' E, 2 , July 2011. **New record for Iran.**

**Distribution outside Iran:** Azerbaijan, Croatia, Czech Republic, France, Germany, Hungary, Moldova, Mongolia, Romania, Serbia, Slovakia, Sweden, Turkey, and Ukraine.

Genus *Torymoides* Walker, 1871

*Torymoides dispar* (Masi, 1916)

**Material examined:** Golestan National Park, Yakhtikalan (1887 m), 2 , June 2012; Mazandaran province, Sari, 36° 30' N 53° 30' E, 1 , April 2013.

**Distribution outside Iran:** Albania, Bulgaria, Croatia, former Czechoslovakia, Hungary, Italy, Macedonia, Moldova, Montenegro, Slovakia, Syria, Transcaucasus, and Turkey.

Genus *Torymus* Dalman, 1820

*Torymus nitidulus* (Walker, 1833)

**Material examined:** Golestan National Park, Dasht-e-Mirza-Bayloo (1589 m), 1 , June 2012. **New record for Iran.**

**Distribution outside Iran:** China, Czech Republic, Germany, Hungary, Mongolia, Netherlands, Romania, Russia, Sweden, Turkey, UK, Ukraine, USA, and former USSR.

## DISCUSSION

In this research, 19 species of Chalcidoidea (including 12 new records) were collected and determined from some regions of the Golestan province, Iran. Golestan province includes a wide spectrum of flora and fauna, the fauna of most insect families have not been studied so far. Therefore, surveys must be continued in order to

determine the fauna of different insect taxa, particularly to find new data on Chalcidoidea. The fauna of Iranian Chalcidoidea is poorly studied and only a few families have been catalogued (e.g. Agaonidae: Ghahari and van Noort, 2011; Aphelinidae: Abd-Rabou *et al.*, 2013; Encyrtidae: Fallahzadeh y Japoshvili, 2010; Eulophidae: Hesami *et al.*, 2011, Talebi *et al.*, 2011; Eurytomidae: Stojanova y Ghahari, 2009). Iran forms a large part of the Iranian plateau, and covers an area of 1,623,779 km<sup>2</sup>, with various geographical regions and climates, so we expect a high diversity of Chalcidoidea in this area of the Palaearctic region.

Many species of Chalcidoidea are efficient and excellent parasitoids of agricultural pests, over 800 species have been associated with biological control programs (Godfray, 1994). Conservation of these natural enemies is necessary in different agroecosystems in order to increase their efficiency for pest control. The inadequate use of pesticides may lead to diminish the chalcidoid population with a consequent explosion of pest populations; beneficial effects of chalcidoids are revealed when the equilibrium is restored (Huffaker and Messenger, 1976; Barbosa, 1998; Bellows and Fisher, 1999).

#### ACKNOWLEDGEMENTS

This research was supported by Islamic Azad University [Yadegar – e- Imam Khomeini (RAH) Branch]. The authors appreciate the help of A. Ribes (University of Lleida, Spain), Z.A. Yefremova (Uljanovsk State Pedagogical University, Russia) and A. Stojanova (National Museum of Natural History, Bulgaria) for the identification of some specimens, and J. S. Noyes (UK) for providing some papers.

#### REFERENCES

ABD-RABOU, SH., H. GHAHARI, S. N. MYARTSEVA, AND E. RUÍZ-CANCINO. 2013. Iranian Aphelinidae (Hymenoptera: Chalcidoidea). *Journal of Entomology and Zoology Studies*, 1(4): 116–140.

BARBOSA, P. 1998. *Conservation biological control*. Academic Press, San Diego, California, USA, 396 p.

BELLOWS, T. S. AND T. W. FISHER. 1999. *Handbook of biological control: principles and applications*. Academic Press, San Diego, New York, 1046 p.

BOUCEK, Z. 1988. Australasian Chalcidoidea (Hymenoptera). A biosystematic revision of genera of fourteen families, with a reclassification of species. CAB International, Wallingford, Oxon, U.K., 832 p.

FALLAHZADEH, M. AND G. JAPOSHVILI. 2010. Checklist of Iranian Encyrtids (Hymenoptera: Chalcidoidea) with descriptions of new species. *Journal of Insect Science*. 10: 1–24.

GAULD, I. D. AND B. BOLTON. 1988. *The Hymenoptera*. Oxford University Press, Oxford, UK, Xi + 332 p.

GHAHARI, H. AND S. VAN NOORT. 2011. A comment on Iranian fig wasps (Chalcidoidea: Agaonidae, Pteromalidae). *Linzer biologische Beiträge*, 43/2: 1247–1252.

GODFRAY, H. C. J. 1994. *Parasitoids, behavioral and evolutionary ecology*. Princeton University Press, USA, 473 p.

GOULET, H. AND J. T. HUBER. 1993. *Hymenoptera of the world: an identification guide to families*. Research Branch, Agriculture Canada, vii + 668 p.

GRISSELL, E. AND M. SCHAUFF. 1997. Chalcidoidea. Pp. 45–116. In: Gibson, G., Huber, J. and Wooley, J. (Eds.). *Annotated keys to the genera of Nearctic Chalcidoidea (Hymenoptera)*. NRC Research Press.

HASSAN Z. B., ZEHZAD, B., FARHANG, B., MAJNOUNIAN, H. AND H. GOSHTASB. 1993. *Golestan National Park*. Department of Environment. Fardin Pub. 203 p.

HESAMI, S., EBRAHIMI, E., OSTOVAN, H. AND Z. YEFREMOVA. 2011. Contribution to the study of Eulophidae (Hymenoptera: Chalcidoidea) of Fars province of Iran: II- Subfamilies Entiinae and Eulophinae, with a preliminary checklist of Eulophidae species in Iran. *Plant Protection Journal*, 2(3): 239–253.

HUFFAKER, C. B. AND P. S. MESSENGER. 1976. *Theory and practice of biological control*. Academic Press, New York, USA, 788 p.

LASALLE, J. AND I. GAULD. 1991. Parasitic Hymenoptera and the biodiversity crisis. *Redia*, 74: 315–334.

NOYES, J. S. 2014. Universal Chalcidoidea database. (Web page: <http://www.nhm.ac.uk/entomology/chalcidoids/index.html>) (Date accessed: May 26th 2015).

NOYES, J. AND M. HAYAT. 1984. A review of the genera of Indo-Pacific Encyrtidae. *Bulletin of the*

*British Museum (Natural History), Entomology Series*, 48(3): 131–395.

NOYES, J. S. AND E. W. VALENTINE. 1989. Chalcidoidea (Insecta: Hymenoptera) - introduction, and review of genera in smaller families. *Fauna of New Zealand*, 18: 1–91.

PECK, O., BOUCEK, Z. AND A. HOFFER. 1964. Keys to the Chalcidoidea of Czechoslovakia (Insecta: Hymenoptera). *Memoirs of the Entomological Society of Canada*, No. 34: 170 p, 289 figs.

STOJANOVA, H. AND H. GHAHARI. 2009. Checklists of Iranian Eurytomidae and Torymidae (Hymenoptera, Chalcidoidea). *Linzer biologische Beiträge*, 41/1: 845–862.

TALEBI, A. A., MOHAMMADI-KHORAMABADI, A. AND E. RAKHSHANI. 2011. Checklist of eulophid wasps (Insecta: Hymenoptera: Eulophidae) of Iran. *Check List*, 7(6): 708–719.